

statins have been widely investigated, and their use is usually recommended in the secondary prevention of acute coronary syndrome (ACS). To investigate the use of statins for the primary prevention of ACS, between April 2004 and February 2005 we enrolled a total of 31 end-stage renal disease patients on MHD, with neither history nor clinical and instrumental signs of earlier major adverse cardiac events (MACEs). We started each patient on atorvastatin (10 mg) daily regardless of serum low-density lipoprotein cholesterol associated with conventional pharmacologic control of blood pressure, serum glucose, and calcium-phosphate balance. During a 4-year follow-up overall 3-year actuarial survival was 67%,³ compared with a 20–40% 5-year survival reported earlier.⁴ Cardiovascular mortality was 12.9%, (4 patients). ‘MACE’ free’ (lethal and non-lethal) 3-year actuarial survival was 53%. Treatment with statins may be considered in the primary prevention of MACEs in hemodialysis patients.

1. Cachofeiro V, Goicochea M, Gracia de Vinuesa S *et al.* Oxidative stress and inflammation, a link between chronic kidney disease and cardiovascular disease. *Kidney Int* 2008; **74**(suppl 111): S4–S9.
2. Maccarrone M, Taccone Gallucci M, Meloni C *et al.* ‘Activation of 5-lipoxygenase and related lipid peroxidation in hemodialysis patients’. *J Am Soc Nephrol* 1999; **10**: 1991–1996.
3. Lonzi M, Manca di Villahermosa S, Summaria F *et al.* Cardiovascular outcome in asymptomatic haemodialysis patients submitted to aggressive medical therapy: a 30-month follow-up. *Nephrol Dial Transplant* 2007; **22**s6: vi322–vi323.
4. Akmal M. Hemodialysis in diabetic patients. *Am J Kidney Dis* 2001; **38**(suppl 1): S195–S199.

Simone Manca-di-Villahermosa¹, Massimo Taccone-Gallucci¹ and Mauro Maccarrone²

¹Department of Internal Medicine, Tor Vergata University, Rome, Italy, and Nephrology and Dialysis Unit, Tor Vergata University Hospital, Rome, Italy and ²Department of Biomedical Sciences, University of Teramo, Italy, and European Center for Brain Research (CERC)/S. Lucia Foundation, Rome, Italy
Correspondence: Massimo Taccone-Gallucci, Unità Operativa di Nefrologia e Dialisi Policlinico Tor Vergata; Viale Oxford 81, Rome 00133, Italy.
E-mail: taccone.gallucci@med.uniroma2.it

Response to ‘Treatment with statins may be considered in ESRD patients for primary prevention of cardiovascular disease’

Kidney International (2009) **75**, 1355; doi:10.1038/ki.2009.111

Considering the clinical benefits of statins for primary and secondary cardiovascular event prevention in normal renal function patients, one can assume a similar benefit for patients with chronic kidney disease. Accordingly, Manca-di-Villahermosa’s letter¹ reports that a small group of end-stage renal disease patients receiving atorvastatin show low cardiovascular mortality after 3 years of follow-up. However, in the 4-D study—the only randomized placebo-controlled statin trial done to date, which included 1255 patients with type 2 diabetes undergoing hemodialysis—atorvastatin administration was not

associated with cardiovascular mortality reduction, which accounts for half the deaths in such patients.² Moreover, this occurs independently of the presence or absence of high c-reactive protein levels that are associated with cardiovascular event risk.³ Reasons for this discrepancy are not evident, as statins exert anti-inflammatory, antioxidant, and lipid-lowering effects as effective as those observed in normal renal function participants. End-stage renal disease patients are at high risk for cardiovascular complications, in whom cardiovascular disease is considered complex and aggravated by coexisting factors, including malnutrition, accelerated atherosclerosis, left ventricular hypertrophy, cardiac fibrosis, and sympathetic overactivity. In the 4-D study, most of cardiovascular deaths were caused by sudden death and not by coronary heart disease.² Data for two ongoing large trials (AURORA and SHARP) would help put into perspective statin’s beneficial effect in chronic kidney disease with respect to cardiovascular mortality. Apart from this, it has been reported that statin use is associated with sepsis incidence reduction—a major cause of morbimortality in chronic kidney disease.⁴ This protective benefit could contribute to total mortality reduction in peritoneal dialysis and hemodialysis patients using statins reported by some observational studies (e.g., DOPPS).⁵

1. Manca-di-Villahermosa S, Maccarrone M, Taccone-Gallucci M. Treatment with statins may be considered in ESRD patients for primary prevention of cardiovascular disease. *Kidney Int* 2009; **75**: 1354–1355.
2. Wanner C, Krane V, März W *et al.* Atorvastatin in patients with type 2 diabetes mellitus undergoing hemodialysis. *N Engl J Med* 2005; **353**: 238–248.
3. Krane V, Winkler K, Drechsler C *et al.* Effect of atorvastatin on inflammation and outcome in patients with type 2 diabetes mellitus on hemodialysis. *Kidney Int* 2008; **74**: 1261–1467.
4. Gupta R, Plantita L, Fink NE *et al.* Statin use and hospitalization for sepsis in patients with chronic kidney disease. *JAMA* 2007; **297**: 1455–1464.
5. Mason NA, Baillie GR, Satyathum S *et al.* HMG-coenzyme A reductase inhibitor use is associated with mortality reduction in hemodialysis patients. *Am J Kidney Dis* 2005; **45**: 119–126.

Victoria Cachofeiro¹, Vicente Lahera¹, Marian Goicochea², Soledad Garcia de Vinuesa² and José Luño²

¹Department of Physiology, School of Medicine, Universidad Complutense, Madrid 28040 and ²Department of Nephrology, Hospital General Universitario Gregorio Marañón, C/ Dr Esquerdo 47, Madrid 28007
Correspondence: Victoria Cachofeiro, School of Medicine, Universidad Complutense, Physiología, Departamento de Fisiología, Facultad de Medicina, UCM, Madrid 28040, Spain. E-mail: vcara@med.ucm.es

On vascular calcification prevention with phosphonoformate and bisphosphonates

Kidney International (2009) **75**, 1355–1356; doi:10.1038/ki.2009.108

To the Editor: Phosphonoformic acid (PFA) is a non-hydrolyzable inorganic pyrophosphate analog, and it has been used as an experimental inhibitor of renal and




Figure 1 | Effect of phosphonoformic acid (PFA) on inorganic phosphate (Pi) transport and calcium deposition in vascular smooth muscle cells.

intestinal inorganic phosphate (Pi) transport.¹ PFA competitively inhibits type II Na/Pi cotransporters (intestinal NaPi-IIb, and renal NaPi-IIa and NaPi-IIc).¹ Conversely, it is not an inhibitor of type III Pi transporters, namely the ubiquitous Pit-1 and Pit-2, as we have shown² and as Professor Giachelli has demonstrated recently in this journal.³

PFA successfully prevents Pi-induced calcification of vascular smooth muscle cells (VSMCs; EC₅₀ of 60 μM),⁴ but it inhibits Pi transport with very low affinity (K_i 2.6 mM)² because these cells only express type III Na/Pi cotransporters.² Therefore, the mechanism of vascular prevention by PFA should be different from that of Pi-transport inhibition, as we have described recently (Figure 1).

In a recent study,⁴ we clarified the mechanism whereby PFA prevents VSMC calcification. We showed that PFA inhibits calcium-phosphate deposition in a process that is independent of any cell activity or metabolism and that is similar to the physicochemical mechanism of bisphosphonate drugs and pyrophosphate.⁴ This rationale was already proposed decades ago by several researchers, and it also coincides with a recent study by Prof. O'Neill's group.⁵ Therefore, the use of PFA as a tool to inhibit Pi transport in vascular calcification research should be avoided, because the concentrations of PFA that significantly inhibit Pi transport in VSMC also exhibit cytotoxic side effects.⁴

1. Villa-Bellosta R, Sorribas V. Role of rat sodium/phosphate cotransporters in the cell membrane transport of arsenate. *Toxicol Appl Pharmacol* 2008; **232**: 125–134.
2. Villa-Bellosta R, Bogaert YE, Levi M *et al*. Characterization of phosphate transport in rat vascular smooth muscle cells: implications for vascular calcification. *Arterioscler Thromb Vasc Biol* 2007; **27**: 1030–1036.
3. Giachelli CM. The emerging role of phosphate in vascular calcification. *Kidney Int* 2009; **75**: 890–897.
4. Villa-Bellosta R, Sorribas V. Phosphonoformic acid prevents vascular smooth muscle cell calcification by inhibiting calcium-phosphate deposition. *Arterioscler Thromb Vasc Biol* 2009; doi:10.1161/ATVBAHA.108.183384 (in press).
5. Lomashvili KA, Monier-Faugere MC, Wang X *et al*. Effect of bisphosphonates on vascular calcification and bone metabolism in experimental renal failure. *Kidney Int* 2009; **75**: 617–625.

Ricardo Villa-Bellosta¹ and Víctor Sorribas¹

¹Laboratory of Molecular Toxicology, University of Zaragoza, Zaragoza, Spain

Correspondence: Ricardo Villa-Bellosta or Víctor Sorribas, Laboratory of Molecular Toxicology, University of Zaragoza. Calle Miguel Servet 177, 50013 Zaragoza, Spain. E-mails: rvilla@unizar.es or sorribas@unizar.es

Response to 'On vascular calcification prevention with phosphonoformate and bisphosphonates'

Kidney International (2009) **75**, 1356; doi:10.1038/ki.2009.113

Despite the fact that phosphonoformic acid (PFA) is a very weak inhibitor of the type III phosphate transporters present in vascular smooth muscle, its inhibition of calcification in cultured vascular smooth muscle cells (VSMCs) has been used to support a role for phosphate transport in vascular calcification.¹ However, as pointed out by Villa-Bellosta and Sorribas,² PFA is also a non-hydrolyzable analog of pyrophosphate (PPi). PPi is a potent, direct inhibitor of hydroxyapatite crystal formation that inhibits vascular calcification *in vitro* and *in vivo*, a property shared by a number of analogs, including bisphosphonates.³ Not surprisingly, Villa-Bellosta and Sorribas have shown that this is also the mechanism by which PFA inhibits calcification in VSMCs. This is yet another example of a 'specific' inhibitor that, like most 'specific' inhibitors, is not specific.

Subsequent studies using antisense RNA directed against Pit-1 have also shown inhibition of calcification in VSMCs,⁴ supporting a role for phosphate transport. However, caution must also be exercised in interpreting these results because VSMCs undergo substantial phenotypic changes in culture and lack the normal elastin matrix, which is the site of medial calcification *in vivo*. Using the whole aorta culture method, we have been unable to duplicate many of the findings in VSMCs related to medial calcification. Thus, the intriguing and potentially important role of phosphate transporters in vascular calcification, although widely cited, remains to be proven in a relevant model.

1. Jono S, McKee MD, Murry CE *et al*. Phosphate regulation of vascular smooth muscle cell calcification. *Circ Res* 2000; **87**: E10–E17.
2. Villa-Bellosta R, Sorribas V. On the vascular calcification prevention with phosphonoformate and bisphosphonates. *Kidney Int* 2009; **75**: 1355–1356.
3. Lomashvili KA, Monier-Faugere MC, Wang X *et al*. Effect of bisphosphonates on vascular calcification and bone metabolism in experimental renal failure. *Kidney Int* 2009; **75**: 617–625.
4. Li X, Yang HY, Giachelli CM. Role of the sodium-dependent phosphate cotransporter, Pit-1, in vascular smooth muscle cell calcification. *Circ Res* 2006; **98**: 905–912.

Koba A. Lomashvili¹ and W. Charles O'Neill¹

¹Emory University, Renal Division, WMB 338, Atlanta, Georgia, USA

Correspondence: W. Charles O'Neill, Emory University, Renal Division, WMB 338, 1639 Pierce Dr, Atlanta, GA 30322, USA.

E-mail: wonell@emory.edu